



## Non-Steroidal Anti-Inflammatory Drugs

(NSAIDs) to Avoid

Donors should not take non-steroidal anti-inflammatory drugs (NSAIDS). Tylenol is the only over the counter pain reliever that should be used post-donation. This is an alphabetical list of common NSAIDS to avoid. This is not an all-inclusive list.

- Actron
- Advil
- Aleve
- Algix
- Anaprox
- Ansaid
- Aspirin
- Acetylsalicylic Acid
- Brufen
- Butazolidin
- Cataflam
- Ceeoxx
- Celebrex
- Celecoxib
- Ceoxx
- Choline Magnesium Trisalicylate
- Clinoril
- Clotam
- Daypro
- Dayrun
- Dexdetoprofen
- Diclofenac
- Diflunisal
- Disalcid
- Dolobid
- Duraprox
- Dynastat
- Etodolac
- Etoricoxib
- Equioxx
- Feldene
- Fenoprofen
- Fenopron
- Firocoxib
- Flurbiprofen
- Flurwood
- Froben
- Ibuprofen
- Indocin, Indocin SR
- Indomethacin
- Keral
- Ketoflam
- Ketoprofen
- Ketorolac
- Licofelone
- Lodine, Lodine XL
- Lornoxicam
- Loxoprofen
- Loxonin
- Loxomac
- Lumiracoxib
- Meclomen
- Meclofenamic Acid
- Meclofenemate
- Medipren
- Mefenamic acid
- Melox
- Meloxicam
- Mesulid
- Midol
- Mobic
- Mobiflex
- Mono-Gesic
- Motrin
- Movalis
- Nabumetone
- Nalfon
- Naprelan
- Naprosyn
- Naproxen
- Nimalox
- Nimesulide
- Nuprin
- Nurofen
- Orudis
- Oruvail
- Oxaporozin
- Oxeno
- Parecoxib
- Phenylbutazone
- Piroxicam
- Ponstel
- Previcox
- Prexige
- Rapid
- Recoxa
- Relafen
- Rofecoxib
- Salflex
- Salicylate
- Salsalate
- Salsitab
- Sprix
- Sulide
- Sulindac
- Tenoxicam
- Tolectin
- Tolfenamic Acid
- Toradol
- Trilisate Disalcid
- Tufnil
- Urbifen
- Valdecoxib (Bextra)
- Vioxx
- Voltaren, Voltaren-XR
- Xefo